

МІНІСТЕРСТВО ОСВІТИ І НАУКИ УКРАЇНИ

ДЕРЖАВНИЙ ВИЩИЙ НАВЧАЛЬНИЙ ЗАКЛАД
«УЖГОРОДСЬКИЙ НАЦІОНАЛЬНИЙ УНІВЕРСИТЕТ»

**НАУКОВИЙ ВІСНИК
УЖГОРОДСЬКОГО УНІВЕРСИТЕТУ**

серія

МАТЕМАТИКА І ІНФОРМАТИКА

Випуск №1 (32)

Ужгород 2018

УДК 51+001

Науковий вісник Ужгородського університету. Сер. матем. і інформ. / Редкол.: В. В. Маринець (гол. ред.) та інші. – Ужгород: Видавництво УжНУ «Говерла», 2018. – випуск №1 (32). – 160 с.

РЕДАКЦІЙНА КОЛЕГІЯ

Головний редактор — Маринець В. В., доктор фізико-математичних наук, професор.
Заст. головн. редактора — Гече Ф. Е., доктор технічних наук, доцент.
Заст. головн. редактора — Король І. І., доктор фізико-математичних наук, доцент.
Відповідальний секретар — Мич І. А., кандидат фізико-математичних наук, доцент.

Члени редакційної колегії:

Бовді А. А., доктор фізико-математичних наук, професор.
Бовді В. А., кандидат фізико-математичних наук, доцент.
Бондаренко В. М., доктор фізико-математичних наук, професор.
Волошин О. Ф., доктор технічних наук, професор.
Головач Й. Г., доктор технічних наук, професор.
Гусак Д. В., доктор фізико-математичних наук, професор.
Задирака В. К., академік НАН України,
доктор фізико-математичних наук, професор.
Козаченко Ю. В., доктор фізико-математичних наук, професор.
Кузка О. І., кандидат фізико-математичних наук, доцент.
Перестюк М. О., академік НАН України,
доктор фізико-математичних наук, професор.
Ронто А. М., доктор фізико-математичних наук, професор,
Ронто М. Й., доктор фізико-математичних наук, професор,
Сливка-Тилищак Г. І., доктор фізико-математичних наук, доцент.
Слюсарчук П. В., кандидат фізико-математичних наук, професор.
Шапочка І. В., кандидат фізико-математичних наук, доцент.

Рекомендовано до друку Вченою радою ДВНЗ «Ужгородський національний університет», протокол № 6 від 21.06.2018 р.

Свідоцтво про державну реєстрацію друкованого засобу масової інформації Серія КВ №7972 від 9.10.2003 року, видане Державним комітетом телебачення і радіомовлення України.

Засновник і видавець – Державний вищий навчальний заклад «Ужгородський національний університет».

Виходить два рази на рік.

Збірник наукових праць видається з 1994 року.

Адреса редакційної колегії: Україна, 88020 Ужгород, вул. Університетська, 14, математичний факультет УжНУ. Тел. (факс): +380 (312) 642725, e-mail: f-mat@uzhnu.edu.ua.

© В. В. Маринець,
І. А. Мич, упорядкування, 2018

© Ужгородський національний університет,
2018

MINISTRY OF EDUCATION AND SCIENCE OF UKRAINE

STATE UNIVERSITY
«UZHHOROD NATIONAL UNIVERSITY»

**SCIENTIFIC BULLETIN OF
UZHHOROD UNIVERSITY**

Series of
MATHEMATICS AND INFORMATICS

Issue no 1 (32)

Uzhhorod 2018

Scientific Bulletin of Uzhhorod University. Series of Mathematics and Informatics / Edit.: V. Marynets (Chief edit.) and others. – Uzhhorod: Scientific Bulletin of UzhNU «Hoverla», 2018. – Issue no 1 (32). – 160 p.

EDITORIAL

Chief editor — Marynets V., Prof., Dr. Sci. (Phys.-Math.).
Deputy Chief editor — Heche F., As. prof., Dr. Sci. (Tech.).
Deputy Chief editor — Korol I., As. prof., Dr. Sci. (Phys.-Math.).
Responsible secretary — Mych I., As. prof., Cand. Sci. (Phys.-Math.).
Members: Bovdi A., Prof., Dr. Sci. (Phys.-Math.).
 Bovdi V., As. prof., Cand. Sci. (Phys.-Math.).
 Bondarenko V., Prof., Dr. Sci. (Phys.-Math.).
 Voloshyn O., Prof., Dr. Sci. (Tech.).
 Holovach J., Prof., Dr. Sci. (Tech.).
 Gusak D., Prof., Dr. Sci. (Phys.-Math.).
 Zadyraka V., Prof., academic of NA of Sc of Ukraine,
 Dr. Sci. (Phys.-Math.).
 Kozachenko Yu., Prof., Dr. Sci. (Phys.-Math.).
 Kuzka A., As. prof., Cand. Sci. (Phys.-Math.).
 Perestyuk N., Prof., academic of NA of Sc of Ukraine,
 Dr. Sci. (Phys.-Math.).
 Ronto A., Prof., Dr. Sci. (Phys.-Math.).
 Ronto M., Prof., Dr. Sci. (Phys.-Math.).
 Slyvka-Tylyshchak G., As. prof., Dr. Sci. (Phys.-Math.).
 Slyusarchuk P., Prof., Cand. Sci. (Phys.-Math.).
 Shapochka I., As. prof., Cand. Sci. (Phys.-Math.).

Recommended for publication by the Scientific Council of UzhNU, record no 6 dated by June 21, 2018.

Certificate of state registration number KV 7972 dated by September 9, 2003.
Founder and Publisher: State University “Uzhhorod National University”.
Published twice a year.
The collection of scientific articles has been published since 1994.

Address of publishing house: Mathematical faculty “UzhNU”, Universytetska str. 14, Uzhhorod, 88020, Ukraine, tel. (fax): +380 (312) 642725, e-mail: f-mat@uzhnu.edu.ua.

ЗМІСТ

1. Андрашко Ю. В., Максим В. В. Булева задача розміщення із урахуванням переваг клієнтів	7
2. Білецька Д. Ю., Шапочка І. В. Тензорні добутки нерозкладних цілочислових матричних зображень симетричної групи третього степеня	15
3. Болдирева В. О., Жмихова Т. В. Ймовірність небанкрутства страхової компанії з витратами на рекламу та інвестиціями у банківський депозит за Каско страхуванням топ-10 страхових компаній України. II.	29
4. Бондаренко В. М., Заціха Я. В. Канонічні форми матричних зображень напівгруп малого порядку	36
5. Бондаренко В. М., Стъопочкіна М. В. Про властивості частково впорядкованих множин ММ-типу (1, 3, 5)	50
6. Брила А. Ю. Про одну задачу лексикографічної оптимізації з інтервальними оцінками	54
7. Глебена М. І., Глебена В. Ф., Попельський О. М. Визначення оптимальних параметрів моделей доступу до інформації у файлах баз даних.	61
8. Дрожжина А. В. Асимптотика розв'язків нелінійних диференціальних рівнянь n -го порядку, що є асимптотично близькими до рівнянь з правильно змінними нелінійностями	67
9. Зубарук О. В. Про зображувальний тип напівгрупи S_{32}^0 над довільним полем	80
10. Кирилюк О. А. Мінімальні незвідні розв'язні підгрупи групи $GL(q, \mathbb{Z}_p)$. . .	86
11. Кічмаренко О. Д. Ступінчасте усереднення керованих функціонально-дифференціальних систем	93
12. Козаченко Ю. В., Василик О. І. Рівномірна збіжність вейвлет-розкладів випадкових процесів з класів $V(\varphi, \psi)$	108
13. Маринець В. В., Питъовка О. Ю. Дослідження крайової задачі для нелінійного хвильового рівняння з розривною правою частиною	116
14. Мич І. А., Ніколенко В. В., Варцаба О. В. Досконалі диз'юнктивні нормальні форми алгебри U_2	124
15. Сапожнікова К. Ю. Часткове усереднення систем диференціальних рівнянь з максимумом	130
16. Сливка-Тилищак Г. І., Михасюк М. М. Властивості узагальненого розв'язку задачі Коші для рівняння теплопровідності на прямій з випадковою правою частиною з простору Орліча	136
17. Чуйко С. М., Чуйко О. С., Чечетенко В. О. Про розв'язання нелінійних інтегрально-диференціальних крайових задач методом Ньютона-Канторовича	147

CONTENTS

1. <i>Andrashko Yu. V., Maksym V. V.</i> Boolean facility location problem with client preferences.....	7
2. <i>Biletska D. Yu., Shapochka I. V.</i> Tensor products of indecomposable integral matrix representations of the symmetric group of third degree.....	15
3. <i>Boldyreva V. O., Zhmykhova T. V.</i> The probability of non-ruin of an insurance company with advertising expenses and investing in bank term deposit by MHull insurance of 10 Top insurance companies of Ukraine. II.....	29
4. <i>Bondarenko V. M., Zaciha Ya. V.</i> Canonical forms of matrix representations of semigroups of small order.....	36
5. <i>Bondarenko V. M., Styopochkina M. V.</i> On properties of posets of MM -type $(1, 3, 5)$	50
6. <i>Bryla A. Yu.</i> On lexicographic optimization problem with interval parameters....	54
7. <i>Hlebena M. I., Hlebena V. F., Popelskyi O. M.</i> Finding the optimum parameters of models of access to information in database files. \\.....	61
8. <i>Drozhzhina A. V.</i> Asymptotic of solutions of the nonlinear differential equations n -th order asymptotically close to the equations with regularly varying nonlinearities.....	67
9. <i>Zubaruk O. V.</i> On representation type of the semigroup S_{32}^0 over an arbitrary field	80
10. <i>Kyryl'uk O. A.</i> Minimal irreducible solvable subgroups of the group $GL(q, \mathbf{Z}_p)$...	86
11. <i>Kichmarengo O. D.</i> Step-by-step averaging of functional-differential control systems.....	93
12. <i>Kozachenko Yu. V., Vasylyk O. I.</i> Uniform convergence of wavelet expansions of random processes from the classes $V(\varphi, \psi)$	108
13. <i>V. V. Marynets, O. Y. Pytovka</i> Investigation of boundary-value problem for non-linear wave equation with discontinuous right part.....	116
14. <i>Mych I. A., Nykolenko V. V., Varcaba E. V.</i> Perfect disjunctive normal forms of algebra U_2	124
15. <i>Sapozhnikova K. Yu.</i> Partial averaging of differential systems with maxima.....	130
16. <i>Slyvka-Tylyshchak G. I., Mykhasiuk M. M.</i> The properties of generalized solution of Cauchy problems for the heat equations with a random right side from Orlicz space.....	136
17. <i>Chuiko S. M., Chuiko O. S., Chechetenko V. O.</i> On of solving nonlinear Noether integral-differential boundary value problems by the of Newton-Kantorovich method.....	147

УДК 517.9

S. M. Chuiko, O. S. Chuiko, V. O. Chechetenko (Donbass State Pedagogical University)

On of solving nonlinear Noether integral-differential boundary value problems by the of Newton-Kantorovich method

Constructive conditions for the existence of a nonlinear Noether integral-differential boundary value problem are found. An iterative scheme with quadratic convergence is constructed to find the solution of a nonlinear integral-differential boundary value problem based on the modification of the Newton-Kantorovich method. In order to justify the quadratic convergence of the modified Newton-Kantorovich method in the case of an undefined system, the original conditions of convergence are proposed.

Знайдені конструктивні умови розв'язності нелінійної нетерової інтегрально-диференціальної крайової задачі. Для розв'язання нелінійної інтегрально-диференціальної крайової задачі на основі методу Ньютона-Канторовича побудовано ітераційну схему з квадратичною збіжністю. Особливістю узагальнення методу Ньютона-Канторовича є його застосування до недовизначених систем.

1. Statement of the problem. We are investigating the problem of constructing a solution [1, 2]

$$y(t) \in \mathbb{D}^2[a; b], \quad y'(t) \in \mathbb{L}^2[a; b]$$

nonlinear Noether ($n \neq p$) integral-differential system

$$y'(t) = A(t)y(t) + \Phi(t) \int_a^b F(y(s), y'(s), s) ds + f(t), \quad (1)$$

that satisfy the boundary condition

$$\ell y(\cdot) = \alpha, \quad \alpha \in \mathbb{R}^p. \quad (2)$$

We seek a solution of the Noetherian boundary value problem (1), (2) in a small neighborhood of solution

$$y_0(t) \in \mathbb{D}^2[a; b], \quad y'_0(t) \in \mathbb{L}^2[a; b]$$

of the generating problem

$$y'_0(t) = A(t)y_0(t) + f(t), \quad \ell y_0(\cdot) = \alpha. \quad (3)$$

Here

$$A(t) \in \mathbb{L}_{n \times n}^2[a; b] := \mathbb{L}^2[a; b] \otimes \mathbb{R}^{n \times n}, \quad \Phi(t) \in \mathbb{L}_{n \times m}^2[a; b], \quad f(t) \in \mathbb{L}^2[a; b];$$

$\ell y(\cdot) : \mathbb{D}^2[a; b] \rightarrow \mathbb{R}^p$ —linear bounded vector functional defined in space $\mathbb{D}^2[a; b]$ n -dimensional absolutely continuous on a segment $[a, b]$ functions. Nonlinear vector-function $F(y(t), y'(t), t)$ twice continuously differentiable in the small neighborhood of the solution $y_0(t)$ generating boundary value problem (3), twice continuously differentiable with respect to $y'_0(t)$, and continuous in the independent variable t

on the segment $[a, b]$. The study of the boundary value problem (1), (2) continues the investigation of the linear integral-differential boundary value problem [2] in the case of a nonlinear integral vector-function $F(y, y', t)$.

2. The generalization of the Newton-Kantorovich theorem. We investigate the problem of finding the solution $z \in \mathbb{R}^n$ of the nonlinear equation

$$\varphi(z) = 0. \quad (4)$$

We assume that the function

$$\varphi(z) : \mathbb{R}^n \rightarrow \mathbb{R}^m, \quad m \neq n$$

is twice continuously differentiable with respect to z in some domain $\Omega \subseteq \mathbb{R}^n$. To construct an iteration scheme $\{z_k\}$, that converges to the solution $\tilde{z} \in \mathbb{R}^n$, we use the Newton method [3–5]. Interest in the use of the Newton-Kantorovich method is associated with its effective application in solving nonlinear equations, as well as in the theory of nonlinear oscillations [3–6], including in the theory of non-linear Noetherian boundary value problems [1, 7, 8].

Suppose an approximation z_k is found that is sufficiently close to an exact solution \tilde{z} of the equation (4). We expand the function $\varphi(z)$ in a neighborhood of the exact solution

$$\varphi(\tilde{z}) = \varphi(z_k) + \varphi'(z_k, \varepsilon) \left(\tilde{z} - z_k \right) + R(\xi_k, \tilde{z} - z_k), \quad (5)$$

where

$$R(\xi_k, \tilde{z} - z_k) := \int_0^1 (1-s) d^2\varphi(\xi_k; \tilde{z} - z_k) ds.$$

Here ξ_k is a point lying between the points \tilde{z} and z_k . In a small neighborhood of the exact solution we have the approximate equality

$$\varphi(z_k) + \varphi'(z_k) \left(\tilde{z} - z_k \right) \approx 0,$$

therefore, in order to find the next approximation of z_{k+1} to the exact solution, it is natural to put

$$\varphi(z_k) + \varphi'(z_k) \left(z_{k+1} - z_k \right) = 0, \quad (6)$$

whence under the condition

$$P_{J_k^*} = 0, \quad J_k := \varphi'(z_k) \in \mathbb{R}^{m \times n} \quad (7)$$

we find

$$z_{k+1} = z_k - J_k^+ \varphi(z_k), \quad k = 0, 1, 2, \dots \quad (8)$$

Here $P_{J_k^*} : \mathbb{R}^m \rightarrow \mathbb{N}(J_k^*)$ is an orthogonal projector of the matrix $J_k^* \in \mathbb{R}^{n \times m}$ and J_k^+ is the pseudoinverse Moore-Penrose matrix [1, 9]. Note that condition (7) is equivalent to the requirement of completeness of the rank matrix J_k and is possible only in case $m \leq n$. We show that the iteration scheme (8) converges to the exact

solution \tilde{z} . Suppose that in the neighborhood of the exact solution \tilde{z} there are inequalities

$$\left\| J_k^+ \right\| \leq \sigma_1(k), \quad \left\| d^2\varphi(\xi_k; \tilde{z} - z_k) \right\| \leq \sigma_2(k) \cdot \|\tilde{z} - z_k\|^2$$

and note that it follows from the equalities (5) and (6) that

$$\varphi'(z_k, \varepsilon) \left(\tilde{z} - z_k \right) = -R(\xi_k, \tilde{z} - z_k),$$

so

$$\|\tilde{z} - z_{k+1}\| \leq \left\| J_k^+ \right\| \cdot \left\| R(\xi_k, \tilde{z} - z_k) \right\| \leq \frac{\sigma_1(k)\sigma_2(k)}{2} \cdot \|\tilde{z} - z_k\|^2.$$

Let there be a constant

$$\theta := \sup_{k \in N} \left\{ \frac{\sigma_1(k)\sigma_2(k)}{2} \right\}.$$

In this case, there is an estimate

$$|\tilde{z} - z_{k+1}| \leq \theta \cdot |\tilde{z} - z_k|^2,$$

which holds that if the iteration scheme (8) converges to the exact solution \tilde{z} of the equation (4), then this convergence is quadratic. Let us find the condition for the convergence of the iteration scheme (8) to the exact solution \tilde{z} of the equation (4). To do this, we make estimates

$$|\tilde{z} - z_1| \leq \theta \cdot |\tilde{z} - z_0|^2, \quad |\tilde{z} - z_2| \leq \theta \cdot |\tilde{z} - z_1|^2 \leq \theta^{1+2} \cdot |\tilde{z} - z_0|^{2^2},$$

$$|\tilde{z} - z_3| \leq \theta \cdot |\tilde{z} - z_2|^2 \leq \theta^{1+2+2^2} \cdot |\tilde{z} - z_0|^{2^3}, \dots,$$

$$|\tilde{z} - z_k| \leq \theta \cdot |\tilde{z} - z_{k-1}|^2 \leq \theta^{1+2+2^2+\dots+2^{k-1}} \cdot |\tilde{z} - z_0|^{2^k}, \dots$$

So there's an inequality [5]

$$|\tilde{z} - z_k| \leq \theta^{\frac{2^k-1}{2-1}} \cdot |\tilde{z} - z_0|^{2^k} = \frac{1}{\theta} \cdot \left(\theta \cdot |\tilde{z} - z_0| \right)^{2^k},$$

indicating the convergence of the iterative process (8) to an exact solution \tilde{z} of the equation (4) under condition

$$\theta \cdot |\tilde{z} - z_0| < 1. \quad (9)$$

In practice, the last inequality can be replaced by the following one:

$$\theta \cdot |z_k - z_0| < 1, \quad k = 1, 2, \dots$$

Lemma. *Suppose that for the equation (4) the following conditions are satisfied.*

- 1) *A non-linear vector-function $f(z) : \mathbb{R}^n \rightarrow \mathbb{R}^m$, twice continuously differentiable with respect to z in some region $\Omega \subseteq \mathbb{R}^n$, in a neighborhood of the point z_0 has a root z^* .*

2) In the neighborhood of the zeroth approximation $z_0 \in \Omega \subseteq \mathbb{R}^n$ there are inequalities

$$\left\| J_k^+ \right\| \leq \sigma_1(k), \quad \left\| d^2\varphi(\xi_k; \tilde{z} - z_k) \right\| \leq \sigma_2(k) \cdot \|\tilde{z} - z_k\|, \quad k = 0, 1, 2, \dots \quad (10)$$

3) The following constant exists

$$\theta := \sup_{k \in N} \left\{ \frac{\sigma_1(k)\sigma_2(k)}{2} \right\}.$$

Then, under conditions (7) and (9), to find the solution z^* of equation (4) the iteration scheme (8) is applicable, and the rate of convergence of the sequence $\{z_k\}$ to the solution z^* of equation (4) is quadratic.

Example 1. The iterative scheme (8) is approximate for finding the solution of the non-linear equation (4), where the vector-function is as follows:

$$\varphi(u) := \begin{pmatrix} x + \sin y + \cos z \\ y + \sin z + \cos x \end{pmatrix}, \quad u := \begin{pmatrix} x \\ y \\ z \end{pmatrix}.$$

This vector-function $\varphi(u) : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ is defined in any open domain $D \subset \mathbb{R}^3$ and is twice continuously differentiable with respect to z in the neighborhood $\Omega \subseteq D \subset \mathbb{R}^3$. We set

$$u_0 := (-0,45 \quad -0,45 \quad -0,45),$$

wherein

$$\text{rank}[\varphi'(u_0)] = 2,$$

besides

$$u_1 \approx (-0,455\,961 \quad -0,457\,894 \quad -0,455\,547)^*,$$

and

$$\text{rank}[\varphi'(u_1)] = 2,$$

Then

$$\left\| [\varphi'(u_1)]^+ \right\|_{\infty} := \sigma_1(1) \approx 2,09\,903, \quad \left\| d^2\varphi(u_1) \right\|_{\infty} := \sigma_2(1) \approx 0,897\,838.$$

In this case, the weakened condition (9)

$$\theta_1 \cdot \|u_1 - u_0\|_{\infty} \approx 0,00743\,856 \ll 1, \quad \theta_1 := \frac{\sigma_1(1)\sigma_2(1)}{2} \approx 0,942\,293$$

is satisfied. Since the condition (9) is satisfied for the first step of the iteration scheme (8), we find

$$u_2 \approx \begin{pmatrix} -0,455\,968\,239\,769\,595 \\ -0,457\,889\,951\,795\,185 \\ -0,455\,537\,594\,550\,856 \end{pmatrix}.$$

Then

$$\text{rank} [\varphi'(u_2)] = 2,$$

besides

$$\left\| [\varphi'(u_2)]^+ \right\|_{\infty} := \sigma_1(2) \approx 2,099, \quad \left\| d^2\varphi(u_2) \right\|_{\infty} := \sigma_2(2) \approx 0,897\,835.$$

In this case, the weakened condition (9)

$$\theta_2 \cdot \|u_2 - u_0\|_{\infty} \approx 0,00743\,453 \ll 1,$$

is satisfied, where

$$\theta_2 := \frac{\sigma_1(2) \sigma_2(2)}{2} \approx 0,00743\,453.$$

For the second step of the iteration scheme (8) the discrepancy of the obtained approximation

$$\|\varphi(u_2)\|_{\infty} \approx 3,69\,679 \times 10^{-11}$$

is sufficiently big, so we find

$$u_3 \approx \begin{pmatrix} 0,455\,968\,239\,730\,150 \\ 0,457\,889\,951\,789\,936 \\ 0,455\,537\,594\,568\,580 \end{pmatrix}.$$

Then

$$\text{rank} [\varphi'(u_3)] = 2,$$

besides

$$\left\| [\varphi'(u_3)]^+ \right\|_{\infty} := \sigma_1(3) \approx 2,099, \quad \left\| d^2\varphi(u_3) \right\|_{\infty} := \sigma_2(3) \approx 0,897\,835.$$

In this case, the weakened condition (9)

$$\theta_3 \cdot \|u_3 - u_0\|_{\infty} \approx 0,00743\,453 \ll 1$$

is satisfied, where

$$\theta_3 := \frac{\sigma_1(3) \sigma_2(3)}{2} \approx 0,942\,278.$$

For the third step of the iteration scheme (8) the discrepancy of the obtained approximation is

$$\|\varphi(u_3)\|_{\infty} \approx 0,$$

so it's natural to confine with this approximation.

The Lemma just proved generalizes the corresponding results [4–8] to the case of matrix J_k irreversibility and can be used in the theory of non-linear Noetherian boundary-value problems [1, 7, 8], in the theory of matrix boundary-value problems [10], and also in the theory of matrix linear differential-algebraic boundary value problem [11–15].

3. Construction of solutions of the integral-differential boundary value problem. Let's denote $X(t)$ normal ($X(a) = I_n$) the fundamental matrix of the homogeneous part of the generating system (3). As you know [1], in a critical case

$$P_{Q^*} \neq 0, \quad Q := \ell X(\cdot) \in \mathbb{R}^{p \times n}$$

generating boundary value problem (3) is solvable if and only if the condition is fulfilled

$$P_{Q_a^*} \left\{ \alpha - \ell K[f(s)](\cdot) \right\} = 0; \quad (11)$$

with r -parametric family of solutions of a problem (3)

$$y_0(t, c_r) = X_r(t)c_r + G[f(s); \alpha](t), \quad c_r \in \mathbb{R}^r$$

depicted by a generalized Green operator

$$G[f(s); \alpha](t) := X(t)Q^+ \left\{ \alpha - \ell K[f(s)](\cdot) \right\} + K[f(s)](t).$$

Here

$$K \left[f(s) \right] (t) := X(t) \int_a^t X^{-1}(s) f(s) ds$$

is Green's operator of the Cauchy problem $y_0(a) = c$ for the generating system (3), $X_r(t) - (n \times r)$ - dimensional matrix, formed from r -linear-independent columns of a normal fundamental matrix $X(t)$; matrix $P_{Q_a^*} \in \mathbb{R}^{d \times p}$, formed from d linear-independent rows of orthoprojector matrix

$$P_{Q^*} : \mathbb{R}^p \rightarrow \mathbb{N}(Q^*).$$

In the critical case, boundary value problem (1), (2) has r -parametric system of solutions

$$y(t) = y_0(t, c_r) + x(t), \quad c_r \in \mathbb{R}^r.$$

To find a deviation

$$x(t) \in \mathbb{D}^2[a; b], \quad x'(t) \in \mathbb{L}^2[a; b]$$

from the generative solution $y_0(t, c_r)$ we get the boundary value problem

$$x'(t) = A(t)x(t) + \Phi(t) \int_a^b F(y(s), y'(s), s) ds, \quad (12)$$

$$\ell x(\cdot) = 0. \quad (13)$$

Deviation

$$x(t) = X(t)v + \Psi(t)u$$

from the generative solution $y_0(t, c_r)$ identify unknown steel

$$u := \int_a^b F(y(s), y'(s), s) ds \in \mathbb{R}^m, \quad v \in \mathbb{R}^m$$

and the matrix

$$\Psi(t) := K[\Phi(s)](t) \in \mathbb{D}_{n \times m}^2[a; b].$$

Solution of the integral-differential system (1) satisfies the boundary condition (2)

$$y(t) = y_0(t, c_r) + x(t), \quad x(t) = X(t)v + \Psi(t)u$$

can be found with the use of the equation

$$Qv + Ru = 0, \quad R := \ell \Psi(\cdot) \in \mathbb{R}^{p \times m}. \quad (14)$$

Let's denote $P_\rho \in \mathbb{R}^{(m+n) \times \rho}$ matrix formed from ρ linearly independent columns of orthoprojector P_0 matrix

$$[Q; R] \in \mathbb{R}^{p \times (m+n)}.$$

Condition (14) satisfying vectors

$$v = P_1 c_\rho, \quad u = P_2 c_\rho, \quad c_\rho \in \mathbb{R}^\rho;$$

here

$$P_0 := \text{col} (P_1, P_2), \quad P_1 \in \mathbb{R}^{n \times \rho}, \quad P_2 \in \mathbb{R}^{m \times \rho}.$$

To find the vector c_ρ , necessary to identify unknowns $u(c_\rho)$ and $v(c_\rho)$ we obtain a nonlinear equation (4); here

$$\begin{aligned} \varphi(c_\rho) := \varphi(u(c_\rho), v(c_\rho)) := & u(c_\rho) - \\ & - \int_a^b F(X(s)v + \Psi(s)u, A(s)X(s)v + \Phi(s)u, s) ds. \end{aligned}$$

If for the obtained equation (4) the conditions of the Lemma are fulfilled, we find the unknown $u(c_\rho)$ та $v(c_\rho)$. Thus, the following theorem is proved.

Theorem. *In the critical case ($P_{Q^*} \neq 0$) generating boundary value problem (3) is solvable if and only if the condition is fulfilled (11); with r -parametric family of solutions of the generating problem (3) looks like*

$$y_0(t, c_r) = X_r(t)c_r + G[f(s); \alpha](t), \quad c_r \in \mathbb{R}^r.$$

The wanted solution

$$y(t) = y_0(t, c_r) + x(t), \quad x(t) = X(t)v + \Psi(t)u, \quad v = P_1 c_\rho, \quad u = P_2 c_\rho, \quad c_\rho \in \mathbb{R}^\rho$$

nonlinear integral-differential boundary value problem (1), (2) defines a vector c_ρ , which satisfies the nonlinear equation (4): $\varphi(c_\rho) = 0$, the conditions of solvability of which is determined by the Lemma. To find the vector c_ρ applicable iterative scheme (8), the rate of convergence of the sequence of approximations to the solution of the equation (4) quadratic.

In the partial case of the equation (4) is linear:

$$\varphi(c_\rho) := \mathcal{B} c_\rho + d, \quad \mathcal{B} \in \mathbb{R}^{m \times \rho},$$

the condition of solvability which is equivalent to demand

$$P_{\mathcal{B}^*} = 0. \quad (15)$$

Here $P_{\mathcal{B}^*} : \mathbb{R}^m \rightarrow \mathbb{N}(\mathcal{B}^*)$ — orthoprojector matrix \mathcal{B}^* . For terms (15) solution of the linear equation (4) looks like

$$c_\rho = P_J c_\mu - J^+ d, \quad c_\mu \in \mathbb{R}^\rho;$$

here $P_J : \mathbb{R}^\rho \rightarrow \mathbb{N}(J)$ — orthoprojector matrix \mathcal{B} .

Example 2. *The requirements of the Theorem satisfy the problem of constructing 2π -periodic solutions of a nonlinear integral-differential system*

$$y' = A(t)y + \Phi(t) \int_0^{2\pi} F(y(s), y'(s), s) ds + f(t); \quad (16)$$

here

$$A := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad f(t) := \begin{pmatrix} \cos t \\ \sin t \end{pmatrix}, \quad \Phi(t) := \begin{pmatrix} \cos^3 t & 0 \\ \cos^3 t & 0 \end{pmatrix},$$

in addition

$$F(y(t), y'(t), t) := y(t)(1 - y^*(t)y'(t)) \cos t.$$

For a generating periodic boundary value problem in the case of a system (16)

$$P_Q = P_{Q^*} = I_2 \neq 0,$$

therefore there is a critical case, and the condition of solvability (11) is fulfilled, with this two-parameter family of solutions of the generating problem

$$y_0(t, c_r) = X_r(t)c_r + G[f(s)](t), \quad c_r \in \mathbb{R}^2$$

depicted by a generalized Green operator

$$G[f(s)](t) = \begin{pmatrix} \sin t + \sin 3t \\ \cos t - \cos 3t \end{pmatrix}$$

and a normal fundamental matrix of the homogeneous part of the generating system (16)

$$X(t) = X_r(t) = \begin{pmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{pmatrix}.$$

Because

$$Q = 0, \quad R = -\frac{3\pi}{4} \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix},$$

get matrices

$$P_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad P_2 = (0 \ 0 \ 1).$$

Periodic solution

$$y(t) = y_0(t, c_r) + x(t), \quad x(t) = X(t)v + \Psi(t)u, \quad v = P_1 c_\rho, \quad u = P_2 c_\rho, \quad c_\rho \in \mathbb{R}^3$$

nonlinear integral-differential system (16) defines a vector c_ρ , which satisfies the linear equation (4):

$$\varphi(c_\rho) := \mathcal{B} c_\rho + d;$$

here

$$\mathcal{B} = \frac{1}{8} \begin{pmatrix} -8\pi & 0 & \pi \\ -\pi & 8 & -8\pi \end{pmatrix},$$

in addition

$$d(c_r) = \frac{\pi}{32} \begin{pmatrix} 1 - 32c_1 + 4c_2 \\ -4(2 + c_1 + 8c_2) \end{pmatrix}, \quad c_r := \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}.$$

The last equation is solvable due to the completeness of the matrix rank \mathcal{B} . Let's put it $c_\mu := 0$; the desired 2π -periodic solution of a nonlinear integral-differential system (16) looks like

$$y(t) = \begin{pmatrix} y_1(t) \\ y_2(t) \end{pmatrix};$$

here

$$\begin{aligned} y_1(t) &= \frac{3\,862\,708 \cos t}{1\,383\,908\,045} + \frac{124\,805 c_1 \cos t}{89\,428\,786} + \frac{12\,394\,757 c_2 \cos t}{1\,110\,179\,952} + \\ &+ \frac{12\,053\,843 \sin t}{539\,822\,396} + \frac{12\,394\,757 c_1 \sin t}{1\,110\,179\,952} + \frac{11\,712\,929 c_2 \sin t}{131\,138\,704} + \frac{\sin 3t}{4}, \\ y_2(t) &= \frac{12\,053\,843 \cos t}{539\,822\,396} + \frac{12\,394\,757 c_1 \cos t}{1\,110\,179\,952} + \frac{11\,712\,929 c_2 \cos t}{131\,138\,704} - \frac{\cos 3t}{4} - \\ &- \frac{3\,862\,708 \sin t}{1\,383\,908\,045} - \frac{124\,805 c_1 \sin t}{89\,428\,786} - \frac{12\,394\,757 c_2 \sin t}{1\,110\,179\,952}. \end{aligned}$$

In a non-critical case ($P_{Q^*} = 0$) solvability condition (11) of integral-differential boundary value problem (1), (2) is satisfied for all inhomogeneities $f(t) \in \mathbb{L}^2[a; b]$ and $\alpha \in \mathbb{R}^p$. This is a typical case for Noether boundary-value problems [1, 2].

Consequence. *In a non-critical case ($P_{Q^*} = 0$) generating problem (3) has an r -parametric family of solutions*

$$y_0(t, c_r) = X_r(t)c_r + G[f(s); \alpha](t), \quad c_r \in \mathbb{R}^r.$$

Then, under condition (14) solutions

$$y(t) = y_0(t, c_r) + x(t), \quad x(t) = X(t)v + \Psi(t)u, \quad v = P_1 c_\rho, \quad u = P_2 c_\rho, \quad c_\rho \in \mathbb{R}^p$$

nonlinear Noether integral-differential boundary value problem (1), (2) defines a vector c_ρ , which satisfies the nonlinear equation (4): $\varphi(c_\rho) = 0$, the conditions of solvability of which is determined by the Lemma. To find the vector c_ρ applicable iterative scheme (8), the rate of convergence of the sequence of approximations to the solution of the equation (4) quadratic.

Example 3. *The requirements of the proved Consequence satisfy the problem of construction 2π -periodic solutions of the scalar integral-differential system*

$$y' = y + \Phi(t) \int_0^{2\pi} F(y(s), y'(s), s) ds + f(t); \quad (17)$$

here

$$f(t) := \sin t, \quad \Phi(t) := \cos t,$$

in addition

$$F(y(t), y'(t), t) := (1 - y^2(t)) (1 - y'^2(t)).$$

For a generating boundary-value problem in the case of a system (17) there is a non-critical case, with the unique solution

$$y_0(t) = G[f(s)](t);$$

here

$$G[f(s)](t) = -\frac{\sin t + \cos t}{2}$$

is generalized Green operator. Periodic solution

$$y(t) = y_0(t) + x(t), \quad x(t) = X(t)v + \Psi(t)u, \quad u = -2v, \quad v \in \mathbb{R}^1$$

nonlinear integral-differential system (17) determines the constant v , which satisfies the nonlinear equation (4):

$$\varphi(v) = -2v - \frac{17\pi}{16} + \frac{7v^2\pi}{2} - \pi v^4.$$

To find a constant v applicable iterative scheme (8), indeed: for terms

$$v_0 := -\frac{1}{2}$$

we get

$$J_0 = -2 - 3\pi \neq 0, \quad v_1 = -\frac{7\pi}{8 + 12\pi},$$

with

$$J_1 = -\frac{256 + 1152\pi + 2512\pi^2 + 3216\pi^3 + 1421\pi^4}{16(2 + 3\pi)^3},$$

$$\sigma_1(1) \approx 0,0894281,$$

in addition

$$\varphi(v_1) \approx 0,0894281.$$

Similarly

$$\sigma_2(1) \approx 13,2612, \quad \theta_1 \approx 0,285343 < 1,$$

hence the condition of convergence of the iterative scheme (8) to the exact solution of the equation (4) the first step is complete. At the second step of the iterative scheme (8) we get

$$v_2 = \frac{\pi}{w} (4352 + 26112\pi + 69728\pi^2 + 91680\pi^3 + 39525\pi^4),$$

with

$$J_2 \approx -11,1795 \neq 0, \quad \sigma_1(2) \approx 0,0894495,$$

in addition

$$\varphi(v_2) \approx 2,70456 \times 10^{-7}.$$

Here

$$w := 16(2 + 3\pi)(256 + 1152\pi + 2512\pi^2 + 3216\pi^3 + 1421\pi^4).$$

Similarly

$$\sigma_2(2) \approx 13,2685, \quad \theta_2 \approx 0,285\,449 < 1,$$

hence the condition of convergence of the iterative scheme (8) to the exact solution of the equation (4) the first step is complete. On the third step of the iterative scheme (8) we get

$$v_3 \approx -\frac{149\,078\,559}{309\,925\,526},$$

with

$$\sigma_1(3) \approx 0,0894\,496, \quad \sigma_2(3) \approx 13,2685.$$

Because

$$\theta_3 \approx 0,285\,449 < 1,$$

condition of convergence of the iterative scheme (8) to the exact solution of the equation (4) the third step is completed; in addition

$$\varphi(v_3) \approx 3,99\,680 \times 10^{-15}.$$

Thus, an approximation to a periodic solution is obtained

$$y(t) \approx -\frac{160\,106\,464 \cos t}{163\,205\,053} - \frac{20\,751\,201 \sin t}{1\,092\,981\,631}$$

integral-differential system (17). To evaluate the accuracy of the approximation found to the periodic solution of the integral-differential system (17) define an deviation

$$\left\| y' - y - \Phi(t) \int_0^{2\pi} F(y(s), y'(s), s) ds - f(t) \right\|_{C[0;2\pi]} \approx 3,44\,169 \times 10^{-15}.$$

In addition, we note the periodicity of the approximation obtained.

In the case of insolubility of a nonlinear integral-differential boundary value problem (1), (2) it can be regularize the same way [16–19]. We also note that the scheme proposed in the paper for the study of a nonlinear integral-differential boundary value problem (1), (2) can be transferred to the integral-differential boundary-value problem with delay [1, 20, 21].

Список використаної літератури

1. *Boichuk A.A., Samoilenko A.M.* Generalized inverse operators and Fredholm boundary-value problems. — Utrecht; Boston: VSP, 2004. — XIV + 317 pp.
2. *Samoilenko A.M., Boichuk A.A., Krivosheya S.A.* Boundary value problems for systems of integro-differential equations with Degenerate Kernel // *Ukrainian Mathematical Journal.* — 1996. — **48.** — № 11. — P. 1785 — 1789.
3. *Bogolyubov N.N., Mitropolsky J.A., Samoilenko A.M.* The method of accelerated convergence in nonlinear mechanics. — Kiev: Scientific thought, 1969. — 248 pp.
4. *Kantorovich L.V., Akilov G.P.* Functional analysis. — Moscow: Nauka. — 1977. — 744 pp.
5. *Dennis J. Schnabel R.* Numerical methods of unconditional optimization and solving nonlinear equations. — Moscow: Mir. — 1988. — 440 pp.
6. *Polyak B.T.* The Newton method and its role in optimization and computational mathematics // *Trudy ICA RAN.* — 2006. — **28.** — P. 48 — 66.

7. *Chuiko S.M., Boichuk I.A., Pirus O.E.* On the approximate solution of an autonomous boundary-value problem the Newton - Kantorovich method // Journal of Mathematical Sciences — 2013. — **189**, № 5. — P. 867 — 881.
8. *Chuiko S.M., Pirus O.E.* On the approximate solution of autonomous boundary-value problems by the Newton method // Journal of Mathematical Sciences — 2013. — **191**, № 3. — P. 449 — 464.
9. *Gantmakher F.R.* Matrix theory. — Moscow: Nauka. — 1988. — 552 pp.
10. *Chuiko S.* Weakly nonlinear boundary value problem for a matrix differential equation // Miskolc Mathematical Notes. — 2016. — **17**, № 1. — P. 139 — 150.
11. *Campbell S.L.* Singular Systems of differential equations. — San Francisco - London - Melbourne: Pitman Advanced Publishing Program. — 1980. — 178 p.
12. *Chuiko S.M.* The Green's operator of a generalized matrix linear differential-algebraic boundary value problem // Siberian Mathematical Journal. — 2015. — **56**, № 4. — pp. 752 — 760.
13. *Boichuk A.A., Pokutnyi A.A., Chistyakov V.F.* Application of perturbation theory to the solvability analysis of differential algebraic equations // Computational Mathematics and Mathematical Physics. — 2013. — **53**. — №6. — P. 777 — 788.
14. *Chuiko S.M.* A generalized matrix differential-algebraic equation // Journal of Mathematical Sciences (N.Y.). — 2015. — **210**, № 1. — P. 9 — 21.
15. *Chuiko S.M.* To the issue of a generalization of the matrix differential-algebraic boundary-value problem // Journal of Mathematical Sciences. — 2017. — **227**, № 1. — P. 16 — 32.
16. *Tikhonov A.N., Arsenin V.Ya* Solution of Ill-Posed Problems. — Winston, Washington, DC. — 1977. — 288 pp.
17. *Krein S.G.* Linear Differential Equations in Banach Space. — Amer. Math. Soc., Providence, RI. — 1971. — 390 pp.
18. *Forghan B., Takook M. Zarei A.* Krein regularization of QED. — Annals of Physics. — 2012. — **327**. — 10.1016/j.aop.2012.06.003.
19. *Chuiko S.M.* On the regularization of a matrix differential-algebraic boundary-value problem // Journal of Mathematical Sciences. — 2017. — **220**, № 5. — P. 591 — 602.
20. *Бігун Я.Й.* Існування розв'язку та усереднення нелінійних багаточастотних задач із запізненням // Укр. мат. журн. — 2007. — **59**, № 4. — С. 435 — 446.
21. *Chuiko S.M., Chuiko A.S.* On the approximate solution of periodic boundary value problems with delay by the least-squares method in the critical case // Nonlinear Oscillations (N.Y.) — **14**. — 2012. № 3. P. 445 — 460.

Одержано 10.01.2018